

Non-Hospital Medical and Surgical
Facilities Accreditation Program

ACCREDITATION STANDARDS

Pre-admission
Evaluation and Selection

Copyright © 2025 by the Non-Hospital Medical and Surgical Facilities Accreditation Program and the College of Physicians and Surgeons of British Columbia.

All rights reserved. No part of this publication may be used, reproduced or transmitted, in any form or by any means electronic, mechanical, photocopying, recording or otherwise, or stored in any retrieval system or any nature, without the prior written permission of the copyright holder, application for which shall be made to:

Non-Hospital Medical and Surgical Facilities Accreditation Program
College of Physicians and Surgeons of British Columbia
300-669 Howe Street
Vancouver BC V6C 0B4

The Non-Hospital Medical and Surgical Facilities Accreditation Program and the College of Physicians and Surgeons of BC has used their best efforts in preparing this publication. As websites are constantly changing, some of the website addresses in this publication may have moved or no longer exist.

Introduction

Patient evaluation is to take place before the day of surgery to allow for pre-operative work-up including necessary testing and consultation(s).

Pre-admission Evaluation and Selection

No.	Description	Reference	Risk	Change
PAE1.0	PRE-ADMISSION EVALUATION AND SELECTION			
PAE1.1	Pre-admission screening ensures all patients booked for surgery are appropriate for admission to a non-hospital facility.			
PAE1.1.1	<p>M Pre-admission screening is conducted by regulated health professionals.</p> <p><i>Intent: The patient's medical record is reviewed by a physician or nurse to ensure the patient is appropriate for the nonhospital setting. There may be indications that the patient should be reviewed by an anesthesiologist prior to the day of surgery (i.e. significant comorbidities, obesity, and obstructive sleep apnea).</i></p>	1, 2	M	
PAE1.1.2	<p>M Pre-admission assessment includes a current physical exam.</p> <p><i>Guidance: The physical exam must include a systems review and full functional inquiry. It may be completed by a family physician, the surgeon or a nurse practitioner and as appropriate, by an oral surgeon, podiatrist or osteopath. Patients with a body mass index (BMI) greater than or equal to 40 must have had the physical exam completed within 60 days of the surgery. For all other patients, the physical exam has been completed within 90 days of the surgery. Physical exams that are over the prescribed number of days must be updated prior to surgery or procedure by the operating surgeon, e.g. changes/no changes dated and signed.</i></p>	2, 3, 4, 5, 6	M	Rev. Guidance
PAE1.1.3	<p>M Pre-admission assessment includes a current medical history.</p> <p><i>Guidance: The medical history must include indication(s) for surgery, comorbidities, previous surgery, medications, allergies and sensitivities. The patient's referring physician may provide many of these elements. Patients with a body mass index (BMI) greater than or equal to 40 must have had their medical history completed within 60 days of the surgery. For all other patients, the medical history has been completed within 90 days of the surgery. Medical histories that are over the prescribed number of days must be updated prior to surgery or procedure by the operating surgeon (e.g. changes/no changes, dated and signed).</i></p>	2, 3, 5, 6, 7, 8, 9	M	Rev. Guidance

No.	Description	Reference	Risk	Change
PAE1.1.4	<p>M Pre-admission assessment includes review of previous anesthetic history. <i>Guidance: The anesthetic history includes the patient's family history of any adverse reactions associated with anesthesia.</i></p>	1, 2, 3, 7, 9, 10	M	
PAE1.1.5	<p>M Pre-admission assessment includes obstructive sleep apnea (OSA) screening using a validated tool (e.g. STOP-Bang), as appropriate. <i>Guidance: The following should be taken into account in determining whether the patient is suitable for the non-hospital setting: OSA severity, coexisting diseases, invasiveness of procedure, type of anesthesia, anticipated post-operative opioid requirements, adequacy of post-discharge observation. All Class 1 and Class 2 facilities are required to screen patients for obstructive sleep apnea (OSA). Class 3 (local anesthesia only) facilities should also screen patients that receive pre-procedural oral sedation for obstructive sleep apnea (OSA).</i></p>	2, 3, 8, 9, 10, 11, 12, 13, 14, 15	M	
PAE1.1.6	<p>M Pre-admission assessment includes venous thromboembolism (VTE) screening. <i>Guidance: Thrombosis Canada provides resources to assist facilities in VTE screening and prophylaxis. The VTE screening is performed by a regulated health professional, preferably the physician as part of the consent discussion, and prior to the day of the procedure. Class 3 (local anesthesia only) facilities are not required to screen patients for VTE. VTE screening is not required for patients undergoing procedures performed using local anesthesia only in Class 1 or Class 2 facilities.</i></p>	2, 7, 9, 16	M	Rev. Guidance
PAE1.1.7	<p>M Pre-admission assessment includes specifying the recommended venous thromboembolism (VTE) prophylaxis based on the patient's VTE score. <i>Guidance: Thrombosis Canada provides resources to assist facilities in VTE screening and prophylaxis. If no prophylaxis is recommended this must be documented. If the planned prophylaxis is different from the prophylaxis recommended by a validated VTE screening tool (e.g. Caprini score), this must be documented with clinical reason(s).</i></p>	7, 9, 16, 17, 18, 19, 20	M	New

No.	Description	Reference	Risk	Change
PAE1.1.8	<p>M Pre-admission assessment includes a patient self-reported questionnaire.</p> <p><i>Guidance: The self-reported questionnaire is an opportunity for the patient to provide information about their medical history, comorbidities, previous surgery, medications, allergies and sensitivities. Scientific literature suggests patient self-reported health information is a valid resource that assists in identifying pre-existing medical conditions that may require further clinical work-up therefore improving the provision of health care and supporting appropriate patient selection.</i></p>	2, 21	M	
PAE1.1.9	<p>M Pre-admission assessment includes infectious diseases and antibiotic resistant organism (ARO) screening.</p> <p><i>Guidance: Screening questions may include: Have you ever been diagnosed (infection or colonization) with an antibiotic-resistant organism (ARO) such as MRSA or VRE? Has anyone in your household ever been diagnosed (infection or colonization) with an ARO such as MRSA or VRE? Have you received health care in a facility outside of Canada in the last 12 months? Have you ever been admitted to, or spent more than 12 continuous hours as a patient in, any health-care facility in the last 12 months?</i></p>	2, 7, 9, 22	M	

No.	Description	Reference	Risk	Change
PAE1.1.10	<p>M Pre-admission assessment includes consultations, as appropriate. <i>Guidance: These include but are not limited to anesthesia, cardiology and internal medicine. In Class 1 (general anesthesia) and Class 2 (IV procedural sedation) facilities, an in-person preoperative anesthetic consultation must be completed before the day of surgery for ASA 3 patients, patients with a BMI greater than or equal to 40 and any patient scheduled for laparoscopic adjustable gastric banding surgery (BMI >30 and <50). An in-person preoperative anesthetic consultation should be completed prior to the day of surgery for patients with a low BMI (less than 18.5), patients with significant comorbidities, and as clinically indicated to ensure the appropriate work-up and consultation(s) are completed. For patients that reside outside of the geographic location of the non-hospital facility where the surgery/procedure will be performed, the in-person anesthetic consultation may be performed by an anesthesiologist where the patient resides (i.e. hospital or non-hospital facility local to the patient). Telemedicine does not satisfy the requirement for an in-person anesthetic consultation due to the physical examination limitations.</i></p>	1, 2, 4, 9	M	
PAE1.1.11	<p>M Pre-admission assessment includes ASA classification. <i>Guidance: The ASA physical classification system is used by physicians (anesthesiologists, surgeons) to predict anesthetic and surgical risk prior to a procedure. The ASA physical classification system guideline provides information on the ASA classifications that may be considered for surgery in the non-hospital setting.</i></p>	1, 2, 4, 9	M	
PAE1.1.12	<p>M Pre-admission assessment includes height, weight and body mass index (BMI). <i>Guidance: Patients are screened using body mass index (BMI). The obesity guideline provides information on the consideration of surgery and anesthesia for patients with elevated BMI.</i></p>	2, 7, 8, 9	M	
PAE1.1.13	<p>M Pre-admission assessment includes preoperative testing based upon the patient's clinical condition(s). <i>Guidance: These include but are not limited to laboratory testing, ECG. The Canadian Anesthesiologists' Society Guidelines refer to Choosing Wisely for recommendations related to preoperative testing</i></p>	1, 2, 4, 7, 9, 21	M	

No.	Description	Reference	Risk	Change
PAE1.11.14	M Pre-admission assessment includes results of radiologic examination, as appropriate.	2, 9	M	
PAE1.11.15	M Pre-admission documentation includes a summative view of sex parameters for clinical use (SPCU) as appropriate. <i>Guidance: In instances where sex assigned at birth may not correctly represent the patient’s sex for health-care purposes and/or where specific additional context is needed to provide safe and affirming care, the pre-admission documentation includes a summative view of SPCU. The SPCU communicates the patient’s “status” by organizing and referring clinicians to artefacts and clinical parameters such as sex-related anatomy (presence or absence of organs), a sex-related medication or hormonal inventory (historically and currently), a surgical inventory, or other useful artefacts necessary to support interpretation of results and clinical decision making. If a summative view does not exist, the summative view will need to be collected and organized such as by talking with the patient about their gender-affirming health care and interventions and by reviewing various progress notes and consult reports.</i>	2, 21, 23	H	
PAE1.2	As part of routine practices, a risk assessment is performed prior to admission. <i>Intent: Patients are to be contacted and a risk assessment performed over the phone within seven days of their scheduled procedure to screen for AROs, illness and skin infection or lesions.</i>			

No.	Description	Reference	Risk	Change
PAE1.2.1	<p>M The risk assessment includes screening for antibiotic resistant organisms (AROs). <i>Guidance: As part of routine practices, a risk assessment is performed prior to admission (i.e. over the phone the day prior to their scheduled procedure) and is documented in the patient's medical record. ARO sample screening questions may include:</i></p> <ul style="list-style-type: none"> • <i>Have you ever been diagnosed (infection or colonization) with an antibiotic-resistant organism (ARO) such as MRSA or VRE?</i> • <i>Has anyone in your household ever been diagnosed (infection or colonization) with an ARO such as MRSA or VRE?</i> • <i>Have you received health care in a facility outside of Canada in the last 12 months?</i> • <i>Have you ever been admitted to, or spent more than 12 continuous hours as a patient in, any health-care facility in the last 12 months?</i> <p><i>The medical director, surgeon and anesthesiologist are to be notified of any positive responses following the risk assessment screening to determine whether it is appropriate to proceed with the surgery as scheduled. Surgery should be cancelled and/or rescheduled if the patient has signs and symptoms of a potentially infectious illness.</i></p>	2, 9, 22	L	
PAE1.2.1	<p>M The risk assessment includes screening for illness. <i>Guidance: As part of routine practices, a risk assessment is performed prior to admission (i.e. over the phone the day prior to their scheduled procedure) and is documented in the patient's medical record. Screening questions for illnesses should cover the following: cough, fever, vomiting, diarrhea, chickenpox (pediatrics), conjunctivitis (ophthalmology). Sample screening question: Have you experienced any of the following symptoms in the last 48 hours: fever, rash, cough, nausea, vomiting, diarrhea? The medical director, surgeon and anesthesiologist are to be notified of any positive responses following the risk assessment screening to determine whether it is appropriate to proceed with the surgery as scheduled. Surgery should be cancelled and/or rescheduled if the patient has signs and symptoms of a potentially infectious illness.</i></p>	9, 22	L	

No.	Description	Reference	Risk	Change
PAE1.3	Policies and procedures contain all of the information necessary for the safety of patients, staff and visitors. <i>Intent: Policies and procedures ensure that activities/procedures are performed consistently and accurately by all personnel within the non-hospital facility</i>			
PAE1.3.1	M There is policy and procedures for pre-admission evaluation and appropriate patient selection.		M	
PAE1.3.2	M There is policy and procedures for patient teaching. <i>Guidance: The preoperative teaching process should begin in the surgeon's office and may continue if a preoperative consultation is needed (i.e. anesthesiology consultation). Patient teaching should include but is not limited to: fasting requirements; preoperative bathing and skin preparation; smoking and alcohol cessation information; taking and holding of medications; items to bring (e.g. CPAP); management of post-operative pain; and review of postoperative instructions including any items required (e.g. crutches). Preoperative teaching is reinforced during admission of the patient for the surgical procedure and post-operative instructions are reviewed with the patient and their family prior to discharge.</i>		M	
PAE1.3.3	M There is policy and procedures for obstructive sleep apnea (OSA) screening and postoperative management. <i>Guidance: OSA is associated with an increased risk of post-operative complications after general anesthesia, neuraxial anesthesia and sedation. OSA screening is an important factor in decreasing the incidence of postoperative complications</i>		M	
PAE1.3.4	M There is policy and procedures for venous thromboembolism (VTE) screening and prophylaxis. <i>Guidance: Thrombosis Canada provides resources to assist facilities in VTE screening and prophylaxis.</i>		M	

References

1. Dobson G, Chau, A, Denomme J, Frost S, Fuda G, McDonnell C, et al. Guidelines to the practice of anesthesia - revised edition 2024. *Can J Anaesth* [Internet]. 2024 [cited 2024 Jul 31];Jan;71(1):8-54.
2. Mackenzie CR, Mandl LA, Reyes C, Lachs M, Magid S. Are ambulatory surgical patients as healthy as we think? Using a self-reported health status questionnaire to identify unsuspected medical comorbidities. *HSS J* [Internet]. 2006 Sep [cited 2024 Jul 31];2(2):121-6.
3. American Society of Anesthesiologists Task Force on Perioperative Management of patients with obstructive sleep apnea. Practice guidelines for the perioperative management of patients with obstructive sleep apnea: an updated report by the American Society of Anesthesiologists Task Force on Perioperative management of patients with obstructive sleep apnea. *Anesthesiology* [Internet]. 2014 Feb [cited 2024 Jul 31]; 120(2):268-86.
4. College of Physicians and Surgeons of Ontario, Out-of-hospital premises inspection program (OHPIP). Out-of-Hospital Premises (OHP) standard: procedures [Internet]. Toronto: College of Physicians and Surgeons of Ontario; 2024 [cited 2024 July 31]. 2 p.
5. College of Physicians and Surgeons of British Columbia, Non-Hospital Medical and Surgical Facilities Accreditation Program. Accreditation standards: infection prevention and control program [Internet]. Version 1.3 Vancouver: College of Physicians and Surgeons of British Columbia; 2024 July 24 [cited 2024 July 31]. 14 p. Document No.: 11729.
6. Medical Council of Canada. Pre-operative medical evaluation [Internet]. Ottawa: Medical Council of Canada; 2017 Feb [cited 2024 Jul 31]. 2 p.
7. Operating Room Nurses Association of Canada (ORNAC). The ORNAC standards, guidelines, and position statements for perioperative registered nurses. 16th ed. Toronto: CSA Group; 2023. 562 p.
8. Joshi GP. Are patients with obstructive sleep apnea syndrome suitable for ambulatory surgery? *ASA Newsletter*. 2006 Jan;70(1);17-9.
9. National Association of PeriAnesthesia Nurses of Canada. Standards for practice. 5th ed. Ottawa, ON: National Association of PeriAnesthesia Nurses of Canada; 2023. 190 p.
10. Subramani Y, Wong J, Nagappa M, Chung F. The benefits of perioperative screening for sleep apnea in surgical patients. *Sleep Med Clin*. 2017 Mar;12(1);123-35.
11. Chung F, Subramanyam R, Liao P, Sasaki E, Shapiro C, Sun Y. High STOP-Bang score indicates a high probability of obstructive sleep apnoea. *Br J Anaesth*. 2012 May;108(5):768-75.

12. Chung F, Abdullah HR, Liao P. STOP-Bang questionnaire: a practical approach to screen for obstructive sleep apnea. *Chest*. 2016 Mar;149(3):631-8.
13. Joshi GP. Patients with obstructive sleep apnea for ambulatory surgery: challenges and management. *Curr Anesthesiol Rep*. 2014 Jun;4:284-89.
14. Swart P, Chung F, Fleetham J. An order-based approach to facilitate postoperative decision-making for patients with sleep apnea. *Can J Anaesth*. 2013 Mar;60(3):321-4.
15. Chung F, Yegneswaran B, Liao P, Chung SA, Vairavanathan S, Islam S, et al. STOP questionnaire: a tool to screen patients for obstructive sleep apnea. *Anesthesiology*. 2008 May;108(5):812-21.
16. Canadian Patient Safety Institute, Safer Healthcare Now! Venous thromboembolism prevention: getting started kit [Internet]. Edmonton: Canadian Patient Safety Institute; 2017 [cited 2024 July 31]. 105 p.
17. Henke PK, Kahn SR, Pannucci CJ, Secemsky EA, Evans NS, Khorana AA, et al. Call to action to prevent venous thromboembolism in hospitalized patients: a policy statement from the American Heart Association. *Circulation*. 2020;141:e914-e931.
18. Thrombosis Canada. Caprini score: risk assessment model for venous thromboembolism and recommended intervention for thromboprophylaxis based on risk of VTE [Internet]. Whitby (ON): Thrombosis Canada; 2018 Apr 30 [cited 2024 Jul 31]. 2 p.
19. Thrombosis Canada. Clinical guides: thromboprophylaxis - non-orthopedic surgery [Internet]. Whitby (ON): Thrombosis Canada; 2024 Nov 15 [cited 2024 Dec 6]. 5 p.
20. Thrombosis Canada. Clinical guides: thromboprophylaxis - orthopedic surgery [Internet]. Whitby (ON): Thrombosis Canada; 2024 Mar 11 [cited 2024 Jul 31]. 5 p.
21. College of Physicians and Surgeons of British Columbia. Practice standard: medical records documentation [Internet]. Version 4.3 Vancouver: College of Physicians and Surgeons of British Columbia; 2022 May [cited 2024 July 31]. 4 p.
22. College of Physicians and Surgeons of British Columbia, Non-Hospital Medical and Surgical Facilities Accreditation Program. Accreditation standards: medical records and documentation [Internet]. Version 6.0 Vancouver: College of Physicians and Surgeons of British Columbia; 2024 Mar 27 [cited 2024 July 31]. 46 p. Document No.: 10829.
23. British Columbia Ministry of Health. British Columbia health information standards: gender, sex and sexual orientation (GSSO) health information standard and guidance [Internet]. Version 3.0. Victoria (BC): British Columbia Ministry of Health; 2023 Mar 29 [cited 2024 July 31]. 94 p.

Revision history

Date	Revisions
March 19 2015	NHMSFAP <i>Pre-admission Evaluation and Selection</i> standard approved (version 1.0) (published April 2015)
December 30, 2017	Bylaws change program name to NHMSFAP (no content changes) (version 1.1)
November 29, 2018	Substantial changes to content and format (version 2.0) (published March 29, 2019)
March 24, 2023	New College logo (no content changes) (version 2.1) (published March 24, 2023)
November 30, 2023	<p>Gender, sex and sexual orientation health information standards and guideline revisions (version 3.0) (effective March 1, 2025)</p> <ul style="list-style-type: none"> • New criterion 1.1.14 pre-admission documentation includes a summative view of sex parameters for clinical use (SPCU) as appropriate • Reference list updated • Risk added
November 27, 2024	ISQuaEEA Logo (no content changes) (version 3.1)
December 6, 2024	<p>Revision (version 4.0) (approved September 12, 2024) (effective within 30 days of notification).</p> <ul style="list-style-type: none"> • New criterion 1.1.7 requires that pre-admission assessment includes specifying the recommended venous thromboembolism (VTE) prophylaxis based on the patient's VTE score. • Section 1.1 numbering adjusted accordingly. <p>Various revisions (version 4.0) (approved November 14, 2024)</p> <ul style="list-style-type: none"> • Revised guidance 1.1.2 about updating physical exam if past prescribed number of days • Revised guidance 1.1.3 about updating medical history if past prescribed number of days.

Date	Revisions
	<ul style="list-style-type: none">• Revised guidance 1.1.6 about VTE screening for procedures performed under local anesthesia only.• Reference list updated.